

Edited by:

**DR. JOSE ERNESTO CONTRERAS PULIDO**

Clinical Research of the

**Hospital Ernesto Contreras R.**

Playas de Tijuana, B.C.N.

1963 -1994

Endowed by:

KEMSA LABORATORIES

Playas de Tijuana, B.C.N.

Mexico

## LIST OF AUTHORS

DR. ERNESTO CONTRERAS RODRIGUEZ, M.D.

Chief Director and Founder of the Centro Medico y Hospital Del Mar

Playas de Tijuana, Baja California Norte, Mexico

DR. ABEL MELLADO PRINCE, M.D.

Medical Director of the Centro Medico y Hospital Del Mar

Playas de Tijuana, Baja California Norte, Mexico

DR. JOSE ERNESTO CONTRERAS PULIDO, M.D

Medical Oncologist and Radiation Oncologist of the Centro Medico y

Hospital de Tijuana, Baja California Norte, Mexico

Medical Director of KEMSA Laboratories

Playas de Tijuana, Baja California Norte, Mexico

DR. SALVADOR RUBIO VELIZ, M.D.

Neurologist of the Centro Medico y Hospital Del Mar

Playas de Tijuana, Baja California Norte, Mexico

DR. LUIS COTA ALVAREZ

Dr. of Veterinary Medicine and Technical Zoologist of KEMSA Labs

Playas de Tijuana, Baja California Norte, Mexico

OFB. PETRA BELTRAN CORONA

Chief of the Production Department of KEMSA Labs

Playas de Tijuana, Baja California Norte, Mexico

QFB. MARTHA NAVARRETE DEL RIO

Chief of the Quality Control Department of KEMSA Labs

Playas de Tijuana, Baja California Norte, Mexico

DR. KANEMATSU SUGIURA, M.D.

Researcher of the Sloan-Kettering Institute for Cancer Research

New York, U.S.A.

DR. HAROLD MANNER

Director of Research of the Department of Biology

Loyola University, Chicago, Illinois

The remaining authors and studies were endowed by the McNaughton Foundation, Sausalito, California.

## **FLUORIDATION-LINKED CANCER**

Studies based upon the U.S. Vital Statistics for fluoridated versus non-fluoridated U.S. cities indicate **a significant** (with a greater than 99% confidence level) **increase in cancer death rates occurring within the first two years of artificial fluoridation.**

The nine organ sites affected and their increase above the normal are:- Mouth, 15%; Oesophagus, 48%; Stomach, 22%; Large Intestine, 31%; Rectum, 51%; Kidney, 10%; Bladder and other urinary organs 22%; other organs specifically female: Breast 15%; Ovary and Fallopian Tube, 15%.

Patients having cancers of these organ sites should be advised that they **should not continue to drink or cook with fluoridated city water** but should substitute bottled spring water or distilled water.

<http://www.worldwithoutcancer.org.uk/fluoridation.html>

Index page: <http://www.worldwithoutcancer.org.uk/analysisindex.html>